

Protect Yourself. Protect Your Baby. Get Tested for Syphilis.

Many people with syphilis do not know they have it.
A simple blood test is the only way to know for sure.



If you are pregnant

Get tested 3 times:

- 1 At your **first prenatal visit**
 - 2 Again during your **third trimester**
 - 3 Again at **delivery**
- **Ask your provider** for a syphilis test.
 - If you test positive, **get treated right away** to help protect your baby.
 - If possible, **encourage partners** to get tested too.



If you are not pregnant

- **Get tested** if you have symptoms, a new partner, multiple partners, or a partner who tested positive.
- **Ask a doctor or clinic** for a syphilis test.
- If you test positive, **get treated and tell your sexual partner(s)** so they can get tested too.
- **Using condoms can reduce the risk**, but condoms do not protect against sores not covered by a condom.



Why it matters

- Syphilis can be **passed to a baby** during pregnancy.
- Without treatment, it can cause serious health problems, **premature birth, low birth weight, pregnancy loss, or even death.**
- The good news: **syphilis is treatable and preventable.**



SCAN FOR
TESTING LOCATIONS

Take the next step


**Ask your provider
about testing today.**

A quick test can give you peace of mind and help protect you, your partner, and your baby.



NEED SUBSTANCE
USE TREATMENT?

**Need help?
Support is available.**
Scan the QR code to find information about substance abuse treatment.



Protect your future. Protect others. Get tested.

A simple test today can make a healthier tomorrow for everyone.